

Adsorption (special ISSAC–9 issue)

Supporting Information

Kinetic models assessment for swelling of coal induced by methane and carbon dioxide sorption

Katarzyna Czerw*, Katarzyna Zarębska, Bronisław Buczek, Paweł Baran

*AGH University of Science and Technology, Faculty of Energy and Fuels,
Aleja Mickiewicza 30, 30-059 Cracow, Poland*




Fig. S1 Application of the Busch equations to the kinetics for (a) CH₄ and (b) CO₂ sorption (Eq. (1)), and the sorption-induced coal swelling kinetics for (c) CH₄ and (d) CO₂ (Eq. (2))




Fig. S2 Application of the Siemons equations to the kinetics for (a) CH_4 and (b) CO_2 sorption (Eq. (3)), and the sorption-induced coal swelling kinetics for (c) CH_4 and (d) CO_2 (Eq. (4))




Fig. S3 Application of the PSOEs to the kinetics for (a) CH₄ and (b) CO₂ sorption (Eq. (7)), and the sorption-induced coal swelling kinetics for (c) CH₄ and (d) CO₂ (Eq. (8))




Fig. S4 Application of the Elovich equations to the kinetics for (a) CH₄ and (b) CO₂ sorption (Eq. (9)), and the sorption-induced coal swelling kinetics for (c) CH₄ and (d) CO₂ (Eq. (10))




Fig. S5 Application of the SE equations to the kinetics for (a) CH₄ and (b) CO₂ sorption (Eq. (11)), and the sorption-induced coal swelling kinetics for (c) CH₄ and (d) CO₂ (Eq. (12))