Psychometric validation of the Japanese version of Alcohol Quality of Life Scale (AQoLS-Japan) in the treatment of patients with alcohol use disorder

Susumu Higuchi MD, PhD, Yoshiya Moriguchi MD, PhD, Kristin Hui Xian Tan, MPH

Corresponding author: Professor Susumu Higuchi, Director, National Hospital Organization Kurihama Medical and Addiction Center, 5-3-1 Nobi, Yokosuka, Kanagawa, 239-0841, Japan h-susumu@db3.so-net.ne.jp

Table e1. AQoLS UK Version

Over the last 4 weeks this has applied to me ...Not at all/A little/Quite a lot/Very much

1	I have felt I miss out on everyday activities with family and friends
2	It has been difficult for me to follow through on plans
3	I have been restricted in the places I can go because of alcohol
4	I have struggled doing physical activities due to alcohol (e.g. DIY, walking, cycling)
5	It has been too much effort to do jobs around the house due to alcohol (e.g. cleaning,
	gardening, maintaining the house)
6	Alcohol has interfered with my ability to work
7	I have cut myself off from other people
8	I have neglected the people close to me
9	Alcohol has damaged my close relationships
10	I have behaved badly towards other people
11	I have felt I miss out on family life because of alcohol
12	I have felt that people have no trust in me
13	Alcohol has interfered with my sex life
14	Alcohol has interfered with my relationships with friends
15	I have struggled to keep on top of my everyday household affairs (e.g. dealing with mail,
	organizing appointments)
16	Alcohol has had a negative effect on my housing situation
17	All my money has gone on alcohol
18	Alcohol has caused me financial difficulties
19	I have felt ashamed of myself
20	I have felt that people look down on me
21	I have felt that I am wasting my life
22	I have worried about the effect alcohol has been having on my health
23	I have worried about alcohol causing problems in my life
24	I have had no appetite
25	I have neglected my appearance
26	I have neglected my general health
27	I have put myself in risky situations
28	I have felt that nothing matters more than alcohol
29	Alcohol has controlled me
30	My life has revolved around alcohol
31	I have planned my days around alcohol
32	I have felt as though I have not been in control of myself
33	I have not had a good night's sleep
34	I have not been getting enough sleep

Table e2 Baseline characteristics by Drinking Risk Level (DRL)

	Low or medium	High or very
	DRL (N=117)	high DRL
		(N=33)
Sex; n (%)		
Male	97 (82.9%)	26 (78.8%)
Female	20 (17.1%)	7 (21.2%)
Age (years); mean (SD)	53.3 (12.7)	51.3 (10.0)
Living status; n (%)		
Alone	44 (37.6%)	14 (42.4%)
Not alone	73 (62.4%)	19 (57.6%)
Employment status; n (%)		
On a job	44 (37.6%)	18 (54.5%)
Working at home	3 (2.6%)	3 (9.1%)
Retired	14 (12.0%)	4 (12.1%)
No job	56 (47.9%)	8 (24.2%)
Current smoker; n (%)	80 (69.0%)	20 (60.6%)
Alcohol use disorder; n (%)		
Abuse	4 (3.4%)	0 (0.0%)
Dependence	113 (96.6%)	33 (100.0%)
Total alcohol consumption (g/day) over 28 days;	15.8 (18.6)	112.5 (44.3)
mean (SD)		
Number of drinking days; mean (SD)	8.4 (10.0)	23.8 (5.0)
Number of heavy drinking days; mean (SD)	4.4 (7.4)	23.1 (5.4)
Age at onset of drinking problem (years); mean	38.4 (13.9)	41.2 (12.6)
(SD)		
Time since diagnosis (years); mean (SD)	4.2 (5.4)	2.3 (5.1)
Number of previous attempts to abstain or reduce	9.9 (21.9)	4.7 (5.9)
alcohol consumption; mean (SD)		
Comorbidities; n (%)		
Cardiovascular	26 (22.2%)	7 (21.2%)
Gastric	24 (20.5%)	4 (12.1%)
Hepatic	49 (41.9%)	15 (45.5%)
Metabolic	23 (19.7%)	4 (12.1%)
Psychiatric	44 (37.6%)	11 (33.3%)

Table e3 Generic quality of life measures at Baseline and follow-up visits

Tubic co (Generic quality of life measures at Ba	Baseline (n=150)	Week 2 (n=145)	Month 3 (n=133)
EQ-5D-3L	I have no problems in walking about	118 (80.3%)	123 (86.6%)	110 (85.9%)
Mobility	I have some problems in walking about	26 (17.7%)		18 (14.1%)
Wiodility	I am confined to bed		17 (12.0%)	
		3 (2.0%)	2 (1.4%)	0 (0.0%)
EO ED 31	Missing (n)	100 (07 00/)	3	5
EQ-5D-3L	I have no problems with self-care	129 (87.2%)	134 (93.1%)	124 (95.4%)
Self-care	I have some problems	19 (12.8%)	10 (6.9%)	5 (3.8%)
	washing/dressing myself	0 (0 00()	0 (0 00()	4 (0.00()
	I am unable to wash or dress myself	0 (0.0%)	0 (0.0%)	1 (0.8%)
	Missing (n)	2	1	3
EQ-5D-3L	I have no problems performing my	110 (74.3%)	114 (79.7%)	111 (84.1%)
Usual	usual activities	_ , , ,,	///	
activities	I have some problems performing my	31 (20.9%)	25 (17.5%)	19 (14.4%)
	usual activities			
	I am unable to perform my usual			
	activities	7 (4.7%)	4 (2.8%)	2 (1.5%)
	Missing (n)	2	2	1
EQ-5D-3L	I have no pain or discomfort	91 (62.3%)	86 (59.7%)	87 (66.9%)
Pain/discomf	I have moderate pain or discomfort	48 (32.9%)	50 (34.7%)	36 (27.7%)
ort	I have extreme pain or discomfort	7 (4.8%)	8 (5.6%)	7 (5.4%)
	Missing (n)	4	1	3
EQ-5D-3L	I am not anxious or depressed	84 (57.1%)	88 (61.1%)	94 (71.8%)
Anxiety/depre	I am moderately anxious or depressed	53 (36.1%)	48 (33.3%)	34 (26.0%)
ssion	I am extremely anxious or depressed	10 (6.8%)	8 (5.6%)	3 (2.3%)
	Missing (n)	3	1	2
Score in VAS	Mean (SD)	68.5 (20.3)	70.4 (21.0)	75.3 (18.7)
	Median (Q1-Q3)	70.0 [60 [.] 0-81 ^{.5}]	75.0 [60 [.] 0-82 [.] 0]	80.0 [70.0-90.0]
	Range	(8.0, 100.0)	(7.0, 100.0)	(10.0, 100.0)
	Missing (n)	9	` 11 ′	6
EQ-5D-3L	Mean (SD)	0.8 (0.2)	0.8 (0.2)	0.8 (0.2)
index value	Median (Q1-Q3)	0.8 [0.7-1.0]	0.8 [0.7-1.0]	0.8 [0.7-1.0]
	Range	(0.1, 1.0)	(0.2, 1.0)	(0.2, 1.0)
	Missing (n)	5	4	7
SF-36:	Mean (SD)	85.7 (16.3)	84.4 (18.2)	86.6 (16.0)
Physical	Median (Q1-Q3)	90.0 [80.0-95.0]	90.0 [75.0-95.0]	90.0 [80.0-100.0]
functioning	Range	(10.0, 100.0)	(15.0, 100.0)	(35.0, 100.0)
	Missing (n)	0	0	3
SF-36: Role	Mean (SD)	73.3 (26.3)	73.0 (25.4)	77.3 (24.5)
limitation due	Median (Q1-Q3)	75.0 [56.3-100.0]	75.0 [50.0-100.0]	87.5 [58.3-100.0]
to physical	Range	(0.0, 100.0)	(0.0, 100.0)	(0.0, 100.0)
health	Missing (n)	0.0, 100.0)	0	3
SF-36: Role	Mean (SD)	72.0 (26.6)	71.7 (28.1)	78.1 (25.6)
limitation due	Median (Q1-Q3)	75.0 [58.3-100.0]	75.0 [50.0-100.0]	91.7 [58.3-100.0]
to emotional	Range	(0.0, 100.0)	(0.0, 100.0)	(0.0, 100.0)
problems	Missing (n)	0.0, 100.0)	0	3
SF-36: Vitality	Mean (SD)	58.4 (20.8)	57.2 (24.0)	64.1 (20.3)
oi -oo. Vitality	Median (Q1-Q3)	56.3 [50.0-75.0]	56.3 [43.8-75.0]	68.8 [50.0-81.3]
	Range			
		(0.0, 100.0)	(0.0, 100.0) 0	(0.0, 100.0) 2
SE 26: Montal	Missing (n)	63 0 (24 7)		
SF-36: Mental	Mean (SD)	63.0 (21.7)	62.6 (22.2)	68.1 (21.3)
health	Median (Q1-Q3)	65.0 [50.0-80.0]	70.0 [45.0-80.0]	70.0 [55.0-85.0]
	Range	(5.0, 100.0)	(0.0, 100.0)	(10.0, 100.0)
05.00.0	Missing (n)	1	0	2
SF-36: Social	Mean (SD)	74.5 (27.2)	73.7 (26.4)	78.5 (26.0)
functioning	Median (Q1-Q3)	75.0 [50.0-100.0]	75.0 [50.0-100.0]	87.5 [62.5-100.0]
	Range	(0.0, 100.0)	(12.5, 100.0)	(0.0, 100.0)
	Missing (n)	1	0	1

SF-36: Pain	Mean (SD)	68.2 (25.1)	67.7 (25.7)	68.8 (26.4)
	Median (Q1-Q3)	72.0 [46.5-92.0]	72.0 [52.0-84.0]	72.0 [52.0-100.0]
	Range	(0.0, 100.0)	(0.0, 100.0)	(10.0, 100.0)
	Missing (n)	1	0	2
SF-36:	Mean (SD)	55.1 (17.4)	54.6 (18.5)	58.0 (20.2)
General	Median (Q1-Q3)	56.0 [45.0-67.0]	57.0 [45.0-67.0]	57.0 [47.0-72.0]
health	Range	(10.0, 100.0)	(0.0, 100.0)	(0.0, 100.0)
	Missing (n)	1	1	3

Table e4. Confirmatory factor analysis: AQoLS interfactor correlations (SE) at Baseline and Week 2

AQoLS Dimension		Activities	Relationships	Living Conditions	Negative Emotions	Control	Sleep	Self- esteem
Activities	Baseline	1.00 (-)						
Relationships	Baseline	0.90 (0.02)	1.00 (-)					
Living Conditions	Baseline	0.80 (0.04)	0.86 (0.04)	1.00 (-)				
Negative Emotions	Baseline	0.84 (0.03)	0.79 (0.04)	0.68 (0.06)	1.00 (-)			
Control	Baseline	0.88 (0.02)	0.84 (0.03)	0.76 (0.05)	0.78 (0.05)	1.00 (-)		
Sleep	Baseline	0.53 (0.06)	0.52 (0.07)	0.51 (0.06)	0.52 (0.08)	0.52 (0.07)	1.00 (-)	
Self-esteem	Baseline	0.84 (0.03)	0.97 (0.02)	0.79 (0.04)	0.85 (0.04)	0.80 (0.04)	0.42 (0.07)	1.00 (-)

アルコール クオリティ・オブ・ライフ スケール (AQoLS)

指示をよく読んでください

本アンケートの目的は、<u>過去4週間に</u>、アルコールが及ぼした影響の程度を調べることです。過去4週間より前のことは回答しないでください。

ここでいう「アルコール」とは、過去4週間におけるアルコールとの全般的な関わりであって、その間飲酒していたかどうかは問いません。

各質問に対し、最も当てはまる回答を選んでください。

過去4週間に当てはまる回答...

	全くない	少し	かなり	非常に
1. 家族や友人との日常的な活動の機会を逸していると ていた	感じ			
2. 予定されていた計画を、最後までやり通すのは困難った	であ			
3. アルコールのせいで、出かけられる場所が限られて	こいた 🔲			
4. アルコールのせいで、身体的な活動が困難であった (例:日曜大工、散歩、運動)				
5. アルコールのせいで、家の周りの事をするのに大変力を要していた(例:掃除、ガーデニング、家のメナンス)				
6. アルコールのせいで、仕事をする能力に支障が出て	こいた 🔲			

	過去4週	間に当	てはまる	回答
	全くない	少し	かなり	非常に
7. 他の人との繋がりを断ち切っていた				
8. 親しい人をおろそかにしていた				
9. アルコールのせいで、身内との人間関係が損なわれていた				
10. 他の人に対して、ひどい態度を取ってしまっていた				
	過去4週	間に当っ	てはまる	回答
	全くない	少し	かなり	非常に
11. アルコールのせいで、家庭生活の機会を逸していると 感じていた				
12. 人に信用されていないと感じていた				
13. アルコールのせいで、性生活に支障が出ていた				
14. アルコールのせいで、友人との関係に支障が出ていた				

	過去4週	過去4週間に当てはまる回答		
	全くない	少し	かなり	非常に
15. 日常の家事を適切にこなし続けることが困難であった (例:郵便物の処理、請求への支払)				
16. アルコールのせいで、住宅事情に悪影響があった (例:立ち退き、実家に転居)				
17. 自分のお金を全て、アルコールに使ってしまった				
18. アルコールのせいで、財政難に陥った				
	過去4週	間に当	てはまる	回答
	過去4週 全くない			
19. 自分のことを恥ずかしく感じていた				
19. 自分のことを恥ずかしく感じていた20. 人に見下されていると感じていた				
20. 人に見下されていると感じていた				

	過去4週	間に当っ	てはまる	回答
	全くない	少し	かなり	非常に
24. 食欲が無かった				
25. 外見をおろそかにしていた				
26. 自分の健康全般をおろそかにしていた				
27. 危険な状況に身を置いていた(例:事故、けんか、治 的トラブルに巻き込まれる)				
	過去4週	間に当っ	てはまる	回答
	過去4週 全くない			回答 非常に
28. アルコールよりも大事なことはないと感じていた				
28. アルコールよりも大事なことはないと感じていた 29. アルコールに支配されていた	全くない	少し		
	全くない	少し		
29. アルコールに支配されていた	全くない	少し		

	過去4週間に当てはまる回答				
	全くない	少し	かなり	非常に	
33. 夜良く眠ることができていなかった					
34. 十分な時間睡眠が取れていなかった					

アンケートにご協力いただきありがとうございます。

最初に戻って、各ページのすべての質問に回答していることを確認して ください。